

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 38/26, C07K 14/605</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/30731 (43) International Publication Date: 24 June 1999 (24.06.99)</p>
<p>(21) International Application Number: PCT/US98/26480 (22) International Filing Date: 14 December 1998 (14.12.98) (30) Priority Data: 60/069,728 16 December 1997 (16.12.97) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HOFFMANN, James, Arthur [US/US]; 4272 Woodland Streams Drive, Greenwood, IN 46143 (US). HERMELING, Ronald, Norbert [US/US]; 915 East Woodhill Drive, Indianapolis, IN 46227 (US). NARASIMHAN, Chakravarthy [IN/US]; 10744 Putnam Place, Carmel, IN 46032 (US). (74) Agents: MACIAK, Ronald, S. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).</p>	<p>(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: GLUCAGON-LIKE PEPTIDE-1 CRYSTALS</p>		
<p>(57) Abstract</p> <p>The invention provides individual tetragonal flat rod shaped or plate-like crystals of glucagon-like peptide-1 related molecules, processes for their preparation, compositions and methods of use. The crystal preparations exhibit extended time action <i>in vivo</i> and are useful for treating diabetes, obesity and related conditions.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

GLUCAGON-LIKE PEPTIDE-1 CRYSTALS**Field of Invention**

The present invention relates to peptide chemistry as it applies to pharmaceutical research and development.

5 The invention provides individual tetragonal flat rod shaped or plate-like crystals of glucagon-like peptide-1 related molecules, processes for their preparation, compositions and uses for these improved crystal forms.

Background of the Invention

10 GLP-1, a 37 amino acid peptide naturally formed by proteolysis of the 160 amino acid precursor protein preproglucagon, was first identified in 1987 as an incretin hormone. GLP-1 is secreted by the L-cells of the intestine
15 in response to food ingestion and has been found to stimulate insulin secretion (insulinotropic action) causing glucose uptake by cells which decreases serum glucose levels (see, e g., Mojsov, S., *Int. J. Peptide Protein Research*, 40:333-343 (1992)). GLP-1 is poorly active. A subsequent
20 endogenous cleavage between the 6th and 7th position produces a more potent biologically active GLP-1(7-37)OH peptide. Approximately 80% of the GLP-1(7-37)OH so produced is amidated at the C-terminal in conjunction with removal of the terminal glycine residue in the L-cells and is commonly
25 referred to GLP-1(7-36)NH₂. Molecules which are reasonably homologous to, or are derived from, or based on these native forms will generally be referred to as GLP's in this specification.

The biological effects and metabolic turnover of
30 the free acid, the amide form, and many of the numerous

-2-

known GLP's are similar and show promise as agents for the treatment of diabetes, obesity and related conditions.

However, many GLP's suffer from extremely short biological half lives, some as short as 3-5 minutes, which makes them unattractive for use as pharmaceutical agents. Presently, the activity of dipeptidyl-peptidase-IV (DPP- IV) is believed to readily inactivate many GLP's and is in part responsible for the very short serum half lives observed. Rapid absorption and clearance following parenteral administration are also factors. Thus, there is a need to find a means for prolonging the action of these promising agents.

One such approach has been to modify these molecules to protect them from *in vivo* cleavage by DPP-IV. For example, see US Pat. No. 5,512,549. In the insulin arts, it has long been known that extended time action can be achieved by administering crystalline protein formulations into the subcutis which act like depots, paying out soluble protein over time.

Heterogeneous micro crystalline clusters of GLP-1(7-37)OH have been grown from saline solutions and examined after crystal soaking treatment with zinc and/or m-cresol (Kim and Haren, *Pharma. Res.* Vol. 12 No. 11 (1995)). Also, crude crystalline suspensions of GLP(7-36)NH₂ containing needle-like crystals and amorphous precipitation have been prepared from phosphate solutions containing zinc or protamine (Pridal, et. al., *International Journal of Pharmaceutics* Vol. 136, pp. 53-59 (1996)). Also, EP 0 619 322 A2 describes the preparation of micro-crystalline forms of GLP-1(7-37)OH by mixing solutions of the protein in pH 7-

-3-

8.5 buffer with certain combinations of salts and low molecular weight polyethylene glycols (PEG). However, such crystalline clusters and crude suspensions are less than ideal for preparing long acting pharmaceutical formulations of GLP's since they are loosely bound heterogeneous clusters of crystals or amorphous-crystalline suspensions which tend to trap impurities and are otherwise difficult to reproducibly manufacture and administer.

Most unexpectedly it was discovered that single tetragonal flat rod shaped or plate-like crystals of various GLP's could be reproducibly formed from a mother liquor containing a GLP dissolved in a buffered solution and a C₁₋₃ alcohol, or optionally a mono or disaccharide, over a wide range of pH conditions. The resulting single flat rod shaped or plate-like crystals are superior to, and offer significant advantages over, the GLP-1(7-37)OH crystal clusters or crude suspensions known in the art.

The single tetragonal flat rod shaped or plate-like crystals of the present invention are less prone to trap impurities and therefore may be produced in greater yields and administered more reproducibly than the known heterogeneous clusters. The crystal compositions of the present invention are pharmaceutically attractive because they are relatively uniform and remain in suspension for a longer period of time than the crystalline clusters or amorphous crystalline suspensions which tend to settle rapidly, aggregate or clump together, clog syringe needles and generally exacerbate unpredictable dosing. Most importantly, the crystal compositions of the present invention display extended, uniform, and reproducible

-4-

pharmacokinetics which can be modulated by adding zinc using conventional crystal soaking techniques or, alternatively, by including zinc in the crystallization solution.

5

Brief Summary of the Invention

The present invention includes processes for preparing single rod-shaped or plate-like crystals of glucagon-like peptide-1 related molecules (GLP's) which comprises preparing a crystallization solution comprising a purified
10 GLP, a buffering agent containing an alcohol or a mono or di saccharide, and optionally, ammonium sulfate or zinc. In another embodiment the GLP crystals having tetragonal flat rod shaped or plate-like morphology selected from the group consisting of a GLP-1 analog, a GLP-1 derivative, a DPP-IV
15 protected GLP, a GLP-1 peptide analog, or a biosynthetic GLP-1 analog are claimed. The invention also includes substantially homogenous compositions of GLP crystals, pharmaceutical formulations and processes for preparing such formulations, and methods for treating diabetes, obesity and
20 related conditions.

Detailed Description of the Invention

By custom in the art, the amino terminus of GLP-1(7-37)OH has been assigned number residue 7 and the
25 carboxy-terminus, number 37. This nomenclature carries over to other GLP's. When not specified, the C-terminal is usually considered to be in the traditional carboxyl form. The amino acid sequence and preparation of GLP-1(7-37)OH is well-known in the art. See US Patent Number 5,120,712, the
30 teachings of which are herein incorporated by reference. For the convenience of the reader the sequence is provided below.

-5-

His-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-
 Tyr-Leu-Glu-Gly-Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-Trp-
 Leu-Val-Lys-Gly-Arg-Gly-COOH (SEQ ID NO:1)

"GLP-1 analog" is defined as a molecule having one
 5 or more amino acid substitutions, deletions, inversions, or
 additions relative to GLP-1(7-37) and may include the d-
 amino acid forms. Numerous GLP-1 analogs are known in the
 art and include, but are not limited to, GLP-1(7-34), GLP-
 1(7-35), GLP-1(7-36)NH₂, Gln⁹-GLP-1(7-37), d-Gln⁹-GLP-1(7-
 10 37), Thr¹⁶-Lys¹⁸-GLP-1(7-37), and Lys¹⁸-GLP-1(7-37), Gly⁸-
 GLP-1(7-36)NH₂, Gly⁸-GLP-1(7-37)OH, Val⁸-GLP-1(7-37)OH,
 Met⁸-GLP-1(7-37)OH, acetyl-Lys⁹-GLP-1(7-37), Thr⁹-GLP-1(7-
 37), D-Thr⁹-GLP-1(7-37), Asn⁹-GLP-1(7-37), D-Asn⁹-GLP-1(7-
 37), Ser²²-Arg²³-Arg²⁴-Gln²⁶-GLP-1(7-37), Arg²³-GLP-1(7-37),
 15 Arg²⁴-GLP-1(7-37), α-methyl-Ala⁸-GLP-1(7-36)NH₂, and Gly⁸-
 Gln²¹-GLP-1(7-37)OH, and the like.

Other GLP-1 analogs consistent with the present
 invention are described by the formula:

R₁- X -Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-Tyr-Leu-
 20 Y-Gly-Gln-Ala-Ala-Lys-Z-Phe-Ile-Ala-Trp-Leu-Val-Lys-
 Gly-Arg-R₂ (SEQ ID NO:2)

wherein: R₁ is selected from the group consisting of L-
 histidine, D-histidine, desamino-histidine, 2-amino-
 histidine, beta-hydroxy-histidine, homohistidine, alpha-
 25 fluoromethyl-histidine, and alpha-methyl-histidine; X is
 selected from the group consisting of Ala, Gly, Val, Thr,
 Met, Ile, and alpha-methyl-Ala; Y is selected from the
 group consisting of Glu, Gln, Ala, Thr, Ser, and Gly; Z is
 selected from the group consisting of Glu, Gln, Ala, Thr,

-6-

Ser, and Gly; and R₂ is selected from the group consisting of NH₂, and Gly-OH.

GLP-1 analogs have also been described in WO 91/11457, and include GLP-1(7-34), GLP-1(7-35), GLP-1(7-36),
5 or GLP-1(7-37), or the amide form thereof, and pharmaceutically-acceptable salts thereof, having at least one modification selected from the group consisting of:

(a) substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine,
10 isoleucine, leucine, methionine, phenylalanine, arginine, or D-lysine for lysine at position 26 and/or position 34; or substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, phenylalanine, lysine, or a
15 D-arginine for arginine at position 36;

(b) substitution of an oxidation-resistant amino acid for tryptophan at position 31;

(c) substitution of at least one of: tyrosine for valine at position 16; lysine for serine at position 18;
20 aspartic acid for glutamic acid at position 21; serine for glycine at position 22; arginine for glutamine at position 23; arginine for alanine at position 24; and glutamine for lysine at position 26; and

(d) substitution of at least one of: glycine, serine,
25 or cysteine for alanine at position 8; aspartic acid, glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, or phenylalanine for glutamic acid at position 9; serine, cysteine, threonine, asparagine, glutamine, tyrosine,
30 alanine, valine, isoleucine, leucine, methionine, or

-7-

phenylalanine for glycine at position 10; and glutamic acid for aspartic acid at position 15; and

(e) substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, or phenylalanine, or the D- or N-acylated or alkylated form of histidine for histidine at position 7; wherein, in the substitutions is (a), (b), (d), and (e), the substituted amino acids can optionally be in the D-form and the amino acids substituted at position 7 can optionally be in the N-acylated or N-alkylated form.

A "GLP-1 derivative" is defined as a molecule having the amino acid sequence of GLP-1(7-37) or of a GLP-1 analog, but additionally having chemical modification of one or more of its amino acid side groups, α -carbon atoms, terminal amino group, or terminal carboxylic acid group. A chemical modification includes, but is not limited to, adding chemical moieties, creating new bonds, and removing chemical moieties. Modifications at amino acid side groups include, without limitation, acylation of lysine ϵ -amino groups, N-alkylation of arginine, histidine, or lysine, alkylation of glutamic or aspartic carboxylic acid groups, and deamidation of glutamine or asparagine. Modifications of the terminal amino include, without limitation, the des-amino, N-lower alkyl, N-di-lower alkyl, and N-acyl modifications. Modifications of the terminal carboxy group include, without limitation, the amide, lower alkyl amide, dialkyl amide, and lower alkyl ester modifications. Lower alkyl is C₁-C₄ alkyl. Furthermore, one or more side groups, or terminal groups, may be protected by protective groups

-8-

known to the ordinarily-skilled protein chemist. The α -carbon of an amino acid may be mono- or dimethylated.

Other GLP-1 derivatives are claimed in U.S. Patent No. 5,188,666, which is expressly incorporated by reference.

5 Such molecules are selected from the group consisting of a peptide having the amino acid sequence:

His-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-Tyr-
 Leu-Glu-Gly-Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-Trp-Leu-
 Val-X (SEQ ID NO:3)

10 and pharmaceutically-acceptable salts thereof, wherein X is selected from the group consisting of Lys-COOH and Lys-Gly-COOH; and a derivative of said peptide, wherein said peptide is selected from the group consisting of: a pharmaceutically-acceptable lower alkyl ester of said
 15 peptide; and a pharmaceutically-acceptable amide of said peptide selected from the group consisting of amide, lower alkyl amide, and lower dialkyl amide.

Yet other GLP-1 derivatives consistent for use in the present invention include compounds claimed in U.S.

20 Patent No. 5,512,549, which is expressly incorporated herein by reference, described by the formula:

R¹-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-Tyr-
 Leu-Glu-Gly-Gln-Ala-Ala-Xaa-Glu-Phe-Ile-Ala-Trp-Leu-
 Val-Lys-Gly-Arg-R³ (SEQ ID NO:4)
 25 |
 R²

wherein R¹ is selected from the group consisting of 4-imidazopropionyl, 4-imidazoacetyl, or 4-imidazo- α , α dimethyl-acetyl; R² is selected from the group consisting of
 30 C₆-C₁₀ unbranched acyl, or is absent; R³ is selected from

-9-

the group consisting of Gly-OH or NH₂; and, Xaa is Lys or Arg, may be used in present invention.

"DPP-IV protected GLP's" refers to GLP-1 analogs which are resistant to the action of DPP-IV. These include
5 analogs having a modified or d amino acid residue in position 8. These also include biosynthetic GLP-1 analogs having Gly or the l amino acid residues Val, Thr, Met, Ser, Cys, or Asp in position 8. Other DPP-IV protected GLP's include des amino His' derivatives.

10 "GLP-1 peptide analogs" are defined as GLP-1 analogs or derivatives which exclude acylated forms.

"Biosynthetic GLP-1 analogs" are defined as any GLP-1 analogs or derivatives which contain only naturally occurring amino acid residues and are thus capable of being
15 expressed by living cells, including recombinant cells and organisms.

"Treating" is defined as the management and care of a patient for the purpose of combating the disease, condition, or disorder and includes the administration of a
20 compound of present invention to prevent the onset of the symptoms or complications, alleviating the symptoms or complications, or eliminating the disease, condition, or disorder. Treating diabetes therefore includes the maintenance of physiologically desirable blood glucose
25 levels in patients in need thereof.

The flat rod shaped or plate-like GLP crystals of the present invention, which are prepared using the claimed process, vary in size and shape to some degree. Generally, they range in size from approximately 2-25 microns (μm) by
30 10-150 μm and are flat, having a depth of approximately 0.5-5 μm . These single crystals form from a single nucleation point and do not appear as multiple spiked star-like clusters known in the art.

-10-

Given the sequence information herein disclosed and the state of the art in solid phase protein synthesis, GLP's can be obtained via chemical synthesis. However, it also is possible to obtain some GLP's by enzymatically
5 fragmenting proglucagon using techniques well known to the artisan. Moreover, well known recombinant DNA techniques may be used to express GLP's consistent with the invention.

The principles of solid phase chemical synthesis of polypeptides are well known in the art and may be found
10 in general texts in the area such as Dugas, H. and Penney, C., Bioorganic Chemistry (1981) Springer-Verlag, New York, pgs. 54-92, Merrifield, J.M., Chem. Soc., 85:2149 (1962), and Stewart and Young, Solid Phase Peptide Synthesis, pp. 24-66, Freeman (San Francisco, 1969).

15 Likewise, the state of the art in molecular biology provides the ordinarily skilled artisan another means by which GLP's can be obtained. Although GLP's may be produced by solid phase peptide synthesis, recombinant methods, or by fragmenting glucagon, recombinant methods are
20 preferable when producing biosynthetic GLP-1 analogs because higher yields are possible.

For purposes of the present invention, GLP-1 peptide analogs and biosynthetic GLP-1 peptide analogs are preferred. More preferred are the DPP-IV protected GLP's,
25 More highly preferred are biosynthetic GLP-1 peptide analogs. Another preferred group of GLP-1 peptide analogs are those which contain a single amino acid substitution at the 8 position which may include d and modified amino acid residues. More highly preferred biosynthetic GLP-1 peptide
30 analogs are those which contain a single amino acid

-11-

substitution at the 8 position, more preferably those which contain Gly or the 1 amino acid residues Val, Thr or Met in the 8 position.

The present invention provides a process for
5 producing individual tetragonal rod shaped GLP crystals from a mother liquor. Under low to neutral pH conditions ranging from about pH 6-7, preferably about 6.4 ± about 0.2, the crystallization solution, or mother liquor, contains a final GLP concentration of about 1-10 mg/ml, preferably 2-7 mg/ml.

10 A number of conventional buffer solutions containing an alcohol or mono or disaccharide are suitable in the practice of the invention. 10 to 50 mM Tris, ammonium acetate, sodium acetate, or Bis-Tris is preferred. The concentration of alcohol ranges from about 2-15% (v/v),
15 preferably 3-13%. Preferred alcohols are selected from the group containing methanol, ethanol, propanol, or glycerol, ethanol being most preferred.

Optionally, the addition of approximately 1% (w/v) ammonium sulfate to the mother liquor will generally
20 increase the yield of crystals. The skilled artisan will also recognize the benefits of adding a preservative such as sodium azide and other such preservatives to the mother liquor to prevent bacterial growth.

In another embodiment, mono or disaccharides may
25 be substituted for the alcohol in the same ratios on a weight to volume basis. Mono or disaccharides suitable for use in the presently claimed process include trehalose, mannitol, glucose, erythrose, ribose, galactose, fructose, maltose, sucrose, and lactose, though trehalose is
30 preferred.

-12-

In yet another embodiment of the present invention, the process may be carried out in a neutral or high pH, zinc-containing environment ranging from about pH 7-10, preferably about pH 7.2-9.7. Under these conditions, the GLP concentration is in the range of approximately 1-20 mg/ml, preferably about 2-10 mg/ml. Total zinc, in a molar ratio to GLP, ranges from about 0.5 - 1.7, preferably 0.6 - 1.5.

Under such neutral or high pH conditions with zinc, suitable buffers and salts range in concentration from about 10-100mM glycine and 0-200mM NaCl, preferably 40-60mM glycine and 0-150 mM NaCl. Preferred buffers are glycine, aspartic acid and Tris. The alcohol or sugar conditions are as stated previously.

Once the mother liquor is prepared, it is allowed to stand at approximately 15-37°C, preferably about 18-25°C, for 12-48 hours until crystallization occurs. The crystals may then be transferred or otherwise handled without any noticeable deleterious effects to the crystalline morphology suggesting that such crystals may be stored for prolonged periods without suffering structural damage.

In another embodiment, a pharmaceutical formulation may be prepared by adding pharmaceutically acceptable excipients, carriers, preservatives, and diluents directly to the mother liquor after the crystals have formed. In this embodiment, crystallization and subsequent additions are performed under sterile conditions. Zinc may be added directly to the mother liquor to effect the incorporation of zinc into the crystals. Preservatives may be added to the mother liquor to provide formulations of crystals suitable

-13-

for multiple injections from the same container. Other excipients, such as antioxidants, buffers, acids and bases for pH adjustments, isotonicity agents and the like, may also be added directly to the mother liquor after the
5 crystals have formed.

In another embodiment, the invention provides homogenous compositions of individual tetragonal flat rod shaped or plate-like crystal of GLP's. Prior to the processes herein disclosed and claimed, such compositions
10 could not be achieved. The compositions of the invention are useful in manufacturing processes and for preparing pharmaceutical formulations having extended time action for the treatment or prevention of diabetes, obesity and related conditions.

The claimed GLP crystals and compositions may optionally be treated with zinc using conventional crystal soaking techniques. By soaking the crystals in about a 0.5 mg/ml solution of zinc, complexes of crystals are formed which serve to extend the time action of the administered
20 GLP. Also, by varying the zinc concentration, the complex composition can be altered leading to longer or shorter time actions.

As noted the invention provides pharmaceutical formulations, which are comprised of single tetragonal flat
25 rod shaped or plate-like crystal of a GLP, together with one or more pharmaceutically acceptable diluents, carriers, or excipients. The crystals can be formulated for parenteral administration for the therapeutic or prophylactic treatment of diabetes, obesity or related conditions. For example,
30 the crystals of the present invention can be admixed with

-14-

conventional pharmaceutical carriers and excipients. The formulations comprising the claimed crystals contain from about 0.5 to 50 mg/ml of the active GLP, and more specifically from about 1.0 to 10 mg/ml. Furthermore, the crystals of the present invention may be administered alone or in combination with other antidiabetic agents. For subcutaneous or intramuscular preparations, a sterile formulation of the crystals of the present invention can be administered as a suspension in the original or modified crystallization mother liquor or in a pharmaceutical diluent such as pyrogen-free distilled water, physiological saline, or 5% glucose solution. A suitable formulation of the crystals of the present invention may be prepared and administered as a suspension in an aqueous base or a pharmaceutically acceptable oil base, e.g., an ester of a long-chain fatty acid such as ethyl oleate.

Pharmaceutically acceptable preservatives such as an alkylparaben, particularly methylparaben, ethylparaben, propylparaben, or butylparaben or chlorobutanol, phenol or meta-cresol are preferably added to the formulation to allow multi-dose use.

The formulation may also contain an isotonicity agent, which is an agent that is tolerated physiologically and imparts a suitable tonicity to the formulation to prevent the net flow of water across the cell membrane. Compounds, such as glycerin, are commonly used for such purposes at known concentrations. Other possible isotonicity agents include salts, e.g., NaCl, dextrose, mannitol, and lactose. Glycerin is the preferred isotonicity agent. The concentration of the isotonicity

-15-

agent is in the range known in the art for parenteral formulations, and for glycerin, is preferably about 16 mg/mL to about 25 mg/mL.

The formulation may also contain a
5 pharmaceutically acceptable buffering agent to control the pH at a desired level. The pH is ideally such as to be acceptable to the patient upon administration, yet one at which the formulation is sufficiently stable, both
10 physically and chemically. Preferably, the pH is controlled from a mildly acidic pH to a mildly basic pH, such as, between about pH 5 and pH 9. More preferably, the pH is between about pH 6 and pH 8. Buffering agents include but are not limited to citrate, acetate, phosphate, Tris, or a
15 basic amino acid, such as, lysine or arginine, which are known to be pharmaceutically acceptable in these pH ranges. Other pharmacologically acceptable buffers for buffering at pH in these ranges are known in the art. The selection and concentration of buffer is well within the skill of the art.

20

Example 1**pH 6.4 with 1.0% Ammonium Sulfate**

12.5 mg of chemically synthesized GLP-1(7-37)OH analog having Val substituted for Ala in position 8 (V8-GLP-1) was weighed into a 3.0 ml glass vial and treated with 2.0
25 ml of 10mM Tris-HCl, 0.02% NaN₃, pH 6.4, to give a clear solution at pH 3.6. The pH of the solution was adjusted to 8.7 with 2N NaOH and then lowered to pH 6.4 with 1N HCl. The solution remained clear during the pH adjustments. The solution was filtered through a 0.22 micron Millex GV13
30 syringe filter (Millipore, Bedford MA) into a new 3.0 ml

-16-

glass vial. The concentration of the V8-GLP-1 stock solution was 4.76 mg/ml as determined from the absorbance at 280 nm and using an extinction coefficient of 2.015 for a 1.0 mg/ml V8-GLP-1 solution in a 1 cm cell. A 0.25 ml aliquot of this V8-GLP-1 stock solution was transferred to a 2.0 ml glass vial. To this solution was added 0.25 ml of a 10mM Tris-HCl, 0.02% NaN₃, pH 6.4, buffer containing 2.0% (NH₄)₂SO₄. The vial was sealed, gently swirled, and then placed at 18 °C. After 36 hours crystalline clusters were identified at 200X magnification. For quantitation, a portion of the mother liquor was removed and centrifuged at 16,000 x g. The V8-GLP-1 content remaining in the clear supernatant was determined from the absorbance at 280 nm as cited above. The crystalline yield was quantitated by subtracting the V8-GLP-1 level in the supernatant from the V8-GLP-1 level in the starting solution. This sample showed a crystallization yield of 63.9%.

Example 2

pH 6.4 with 1% Ethanol and 1.0% Ammonium Sulfate

A 0.25 ml aliquot of the V8-GLP-1 stock solution was transferred to a 2.0 ml glass vial as in Example 1. To this solution was added 0.25 ml of a 10mM Tris-HCl, 0.02% NaN₃, pH 6.4, buffer containing 2.0% (NH₄)₂SO₄ and 2.0% ethanol. The solution was then treated and evaluated as in Example 1. This sample generated crystalline clusters and a few single tetragonal crystals. The yield was 73.1%.

-17-

Example 3**pH 6.4 with 5% Ethanol and 1.0% Ammonium Sulfate**

A 0.25 ml aliquot of the V8-GLP-1 stock solution was transferred to a 2.0 ml glass vial as in Example 1. To this solution was added 0.25 ml of a 10mM Tris-HCl, 0.02% NaN₃, pH 6.4, buffer containing 2.0% (NH₄)₂SO₄ and 10.0% ethanol. The solution was then treated and evaluated as in Example 1. This sample generated crystalline clusters, single tetragonal crystals, and some rods. The yield was 80.3%.

Example 4**pH 6.4 with 10% Ethanol and 1.0% Ammonium Sulfate**

A 0.25 ml aliquot of the V8-GLP-1 stock solution was transferred to a 2.0 ml glass vial as in Example 1. To this solution was added 0.25 ml of a 10mM Tris-HCl, 0.02% NaN₃, pH 6.4, buffer containing 2.0% (NH₄)₂SO₄ and 20.0% ethanol. The solution was then treated and evaluated as in Example 1. This sample generated single tetragonal crystals and rods. The yield was 81.9%.

Example 5**pH 6.4 with 1% Ethanol**

A 0.25 ml aliquot of the V8-GLP-1 stock solution was transferred to a 2.0 ml glass vial as in Example 1. To this solution was added 0.25 ml of a 10mM Tris-HCl, 0.02% NaN₃, pH 6.4, buffer containing 2.0% ethanol. The solution was then treated and evaluated as in Example 1. This sample generated a trace of crystal clusters. The yield was 8.8%.

30

-18-

Example 6**pH 6.4 with 5% Ethanol**

A 0.25 ml aliquot of the V8-GLP-1 stock solution was transferred to a 2.0 ml glass vial as in Example 1. To this solution was added 0.25 ml of a 10mM Tris-HCl, 0.02% NaN₃, pH 6.4, buffer containing 10.0% ethanol. The solution was then treated and evaluated as in Example 1. This sample generated crystal clusters, single tetragonal crystals, and rods. The yield was 39.1%.

10

Example 7**pH 6.4 with 10% Ethanol**

A 0.25 ml aliquot of the V8-GLP-1 stock solution was transferred to a 2.0 ml glass vial as in Example 1. To this solution was added 0.25 ml of a 10mM Tris-HCl, 0.02% NaN₃, pH 6.4, buffer containing 20.0% ethanol. The solution was then treated and evaluated as in Example 1. This sample generated single tetragonal crystals and rods. The yield was 55.5%.

20

Example 8**Pharmacokinetics**

28 mg of biosynthetic V8-GLP-1 was weighed into a glass vial and dispersed in 4.5 ml of 10mM NH₄OAc to give a turbid solution with a pH of 5.6. The material was completely solublized by adjusting the pH to 9.5 with 5N NaOH and remained completely soluble after the pH of the solution was lowered to 6.4 with 2N HCl. This solution was filtered through a 0.22 micron Millex GV13 syringe filter into a new glass vial to give a total volume of 4.3 ml. The

30

-19-

concentration of the V8-GLP-1 solution was 5.51 mg/ml as determined from the absorbance at 280 nm of a 20X dilution of the stock solution and using an extinction coefficient of 2.015 for a 1 mg/ml V8-GLP-1 solution in a 1 cm cell. To this solution was added 4.3 ml of a 10mM NH_4OAc , 2.0% $(\text{NH}_4)_2\text{SO}_4$, 20% ethanol, pH 6.4, precipitant buffer. The vial was sealed, the solution was gently swirled and then placed at 18 °C. After 72 hours single tetragonal crystals were identified at 200X magnification. The crystals were removed from the mother liquor by low speed centrifugation and resuspended in a 10 mM NH_4OAc , 16 mg/ml glycerin, pH 5.5, buffer (buffer A) to a concentration of about 4.0 mg/ml. A portion of the mother liquor was centrifuged at 16,000 x g. The V8-GLP-1 content remaining in the clear supernatant was determined from the absorbance at 280 nm. The crystallization yield was quantitated by subtracting the V8-GLP-1 level in the supernatant from the V8-GLP-1 level in the starting solution. This crystallization gave an 83% yield.

Calculated aliquots of 4.0 mg/ml V8-GLP-1 crystal suspensions prepared in a similar manner as above were transferred to five glass vials and diluted with buffer A to a concentration slightly above the final target concentration of 2.5 mg/ml V8-GLP-1. To the crystalline suspensions were added aliquots of a ZnCl_2 stock solution (33.4 mg/ml Zn^{++} in buffer A) to make final zinc concentrations either 0.5, 1.0, 1.5, or 2.4 mg/ml zinc. The suspensions were gently swirled and placed at 5 °C for 18 hours. The final V8-GLP-1 concentration in each vial was now at the 2.5 mg/ml target concentration. After 18 hours

-20-

the crystalline V8-GLP-1 zinc suspensions were transferred to room temperature, passed through a 30 gauge needle, and adjusted to pH 6.0 with 1N NaOH.

A 0.1 mg/ml zinc crystalline V8-GLP-1 suspension was prepared by first treating a 2.5 mg/ml crystal suspension with 0.15 mg/ml zinc in the same manner as described above. After 18 hours at 5 °C the zinc treated crystals were isolated by low speed centrifugation and transferred to buffer B (buffer A containing 0.1 mg/ml zinc). The final V8-GLP-1 concentration of this suspension was adjusted to the 2.5 mg/ml target concentration using buffer B. The suspension was passed through a 30 gauge needle, and the pH increased to 6.0 with 1N NaOH.

The five crystalline V8-GLP-1 zinc suspensions described above, each at 2.5 mg/ml V8-GLP-1 and containing 0.1, 0.5, 1.0, 1.5, or 2.4 mg/ml zinc were tested in overnight-fasted beagle dogs. Each animal received a single 24 nmole/kg subcutaneous injection of the crystalline V8-GLP-1 zinc suspension at time zero. Arterial blood samples (1.5 ml) were withdrawn from the animals at scheduled times, transferred to tubes pretreated with EDTA and containing 40 ul of Trasylol, and then centrifuged. The plasma portion of each sample was separated and stored at -80 °C until analysis. The plasma concentration of immunoreactive V8-GLP-1 in each sample was measured using a RIA procedure. Table 1 shows the resulting immunoreactive V8-GLP-1 plasma levels over a 24 hour time period for each suspension.

-21-

Table 1:

Immunoreactive V8-GLP-1
Levels (picomolar) in Dog Plasma.

Time (hrs)	0.1 mg/ml Zinc (n=5)		0.5 mg/ml Zinc (n=5)		1.0 mg/ml Zinc (n=3)		1.5 mg/ml Zinc (n=5)		2.4 mg/ml Zinc (n=5)	
	V8-GLP-1 (pM)	SEM	V8-GLP-1 (pM)	SEM	V8-GLP-1 (pM)	SEM	V8-GLP-1 (pM)	SEM	V8-GLP-1 (pM)	SEM
0	0	0	0	0	0	0	0	0	0	0
1.5	nd	nd	123	41	27	4	7	5	5	5
3.0	nd	nd	132	38	38	7	40	13	27	21
4.5	nd	nd	196	51	108	41	76	34	83	37
6.0	301	57	264	79	140	55	142	47	143	60
7.5	nd	nd	265	71	184	57	198	61	179	63
9.0	nd	nd	344	94	220	61	252	64	214	66
10.5	nd	nd	302	80	231	78	250	66	225	50
12.0	nd	nd	282	78	236	76	267	60	238	42
13.5	nd	nd	238	54	241	97	286	74	236	38
15.0	nd	nd	263	67	273	118	325	114	246	28
16.5	nd	nd	235	51	234	106	275	77	218	25
18.0	nd	nd	210	47	184	62	254	59	211	23
19.5	nd	nd	221	54	209	120	278	57	173	9
21.0	nd	nd	215	54	219	115	301	48	178	13
22.5	nd	nd	224	54	193	72	232	23	167	9
24.0	190	30	210	51	187	72	227	34	166	25
30.0	nd	nd	nd	nd	nd	nd	nd	nd	171	24

5

SEM = Standard Error of Mean.

nd = not determined

-22-

Example 9**pH 9.4 with 5% Trehalose and Zinc**

6.8 mg of lyophilized biosynthetic V8-GLP-1 was weighed into a 3.0 ml glass vial. Then, 1.0 ml of a 25 mM glycine-HCl, 150 mM NaCl, 5% trehalose, pH 9.0, buffer was added to dissolve the peptide. The solution was then adjusted to pH 10.3 with 5N NaOH. While the solution was gently stirred 9.0 ul of a 10 mg/ml zinc chloride solution in water was added and the pH adjusted to 9.4 with 2N HCl. The final concentration of V8-GLP-1 was 5.4 mg/ml as determined from the absorbance at 280 nm of a 10X dilution of the solution. The solution was then filtered with a 0.22 micron Millex GV13 syringe filter. The vial was capped, gently swirled, and then placed at ambient temperature. After 24 hours V8-GLP-1 crystal clusters and single rectangular crystals were identified at 430X magnification and estimated to be about 40 microns long, 15 microns wide, and 3 microns thick. A portion of the mother liquor was removed and centrifuged at 16,000 x g. The V8-GLP-1 content remaining in the clear supernatant was determined from the absorbance at 280 nm. The crystalline yield was quantitated by subtracting the V8-GLP-1 level in the supernatant from the V8-GLP-1 level in the starting solution. This sample showed a crystallization yield of 89.8%. The small rectangular crystal morphology was not observed in crystallization trials without trehalose.

-23-

Example 10**pH 9.4 with 10% Mannitol and Zinc**

6.8 mg of lyophilized biosynthetic V8-GLP-1 was weighed into a 3.0 ml glass vial, treated with 1.0 ml of a 25 mM glycine-HCl, 150 mM NaCl, 10% mannitol, pH 9.0, buffer and dispersed to give a clear solution. The solution was then adjusted to pH 10.3 with 5N NaOH. While the solution was gently stirred 9.0 ul of 10 mg/ml zinc chloride solution in water was added and the pH adjusted to 9.4 with 2N HCl. The final concentration of V8-GLP-1 was 5.31 mg/ml as determined from the absorbance at 280 nm of a 10X dilution of the crystallization solution. The solution was then filtered with a 0.22 micron Millex GV13 syringe filter. The vial was capped, gently swirled, and then placed at ambient temperature. After 24 hours, small rectangular plate-like crystals of V8-GLP-1 were identified at 430X magnification and estimated to be about 10 to 30 microns long and 10 microns wide. The yield was determined as in Example 9. This sample showed a crystallization yield of 35%.

20

Example 11**pH 9.0 with Zinc**

A 1-ml aliquot of a solution of V8-GLP-1 at 3 mg/ml in 50 mM glycine-150 mM NaCl buffer at pH 9.0 was prepared. To this solution was added 7.5 μ l of a 20.85 mg/ml zinc chloride solution in water, followed by a pH adjustment back up to pH 9.0. After gentle swirling, the clear sample in a 3-ml glass vial was stored at ambient temperature for one day. After this time the crystalline precipitate was examined under the microscope at 90X

30

-24-

magnification, revealing clusters of small plates. For quantitation of crystallization yield, the entire suspension was passed through a 0.2 μ m filter (Gelman Sciences, Ann Arbor, Michigan). The V8-GLP-1 content remaining in the clear filtrate was quantitated by spectroscopic evaluation at a wavelength of 280 nm, using an extinction coefficient of 2.015 for a 1 mg/ml solution of V8-GLP-1 in a 1 cm cell. The crystallization yield was quantitated by subtracting the V8-GLP-1 level in the supernatant from the V8-GLP-1 level in the starting solution. This sample showed a crystallization yield of 5.6%.

Example 12

pH 9.0 with 10% Ethanol and Zinc

A 1-ml aliquot of a solution of V8-GLP-1 was prepared as in Example 11, except that 110 μ l of absolute ethanol was added to the solution prior to the addition of the zinc chloride solution. This sample generated large tetragonal crystals, with some clusters, in 80.6 % yield.

Example 13

pH 9.5 with 10% Ethanol and Zinc

A solution of V8-GLP-1 at 10 mg/ml in 50 mM glycine-150 mM NaCl buffer at pH 10.5 was passed through a sterile 0.2 μ m Acrodisc filter (Gelman Sciences, Ann Arbor, Michigan). To 500 μ l of this solution was added 500 μ l of a 50 mM glycine-150 mM NaCl buffer at pH 9.0. To this solution was then added 110 μ l of absolute ethanol followed by 7.5 μ l of a 20.85 mg/ml zinc chloride solution in water. Small additions of 1N HCl were used to adjust the solution

-25-

to pH 9.5. After gentle swirling the final solution was enclosed in a 3-ml glass vial and stored at ambient temperature for two days. Individual crystalline plates of V8-GLP-1 up to 150 μm in length, approximately 25 μm wide and less than 5 μm thick were generated in 72% yield.

Example 14

pH 7.9 with 8.5% Ethanol and Zinc

V8-GLP-1 was prepared at 4 mg/ml in 50 mM glycine pH 9.5 buffer, followed by passage through a 0.2 μm filter (Gelman Sciences, Ann Arbor, Michigan). To 1-ml of this solution was added 100 μl of absolute ethanol and then 60 μl of 2.08 mg/ml zinc chloride solution in water. Small additions of 0.1N HCl were used to adjust the solution to pH 8.0. After gentle swirling the final solution was enclosed in a 3-ml glass vial and stored at ambient temperature for two hours. The pH of the clear solution was then adjusted to pH 7.86 with small additions of 0.1N HCl and storage at ambient temperature continued for two days. Microscopic examination revealed modest-sized, individual tetragonal plates and some clusters. The V8-GLP-1 content remaining in the clear supernatant was quantitated by spectroscopic evaluation at a wavelength of 280 nm, using an extinction coefficient of 2.015 for a 1 mg/ml solution of V8-GLP-1 in a 1 cm cell. The crystallization yield was quantitated by subtracting the V8-GLP-1 level in the supernatant from the V8-GLP-1 level in the starting solution. This sample showed a crystallization yield of 92.2%.

-26-

Example 15**pH 8.3 with 10% Ethanol and Zinc**

V8-GLP-1 was prepared at 7 mg/ml in 100 mM glycine
pH 10.5 buffer, followed by passage through a 0.2 μ m filter
5 (Gelman Sciences, Ann Arbor, Michigan). To 0.5 ml of this
solution was added 0.4 ml of water. Then 100 μ l of absolute
ethanol was added, followed by about 6 μ l of a 20.86 mg/ml
zinc chloride solution in water. Small additions of 1N HCl
10 were used to adjust the solution to pH 8.33. After gentle
swirling the final solution was enclosed in a 3-ml glass
vial and stored at ambient temperature for one day.
Microscopic examination revealed small, individual
tetragonal plates and some clusters. The V8-GLP-1 content
remaining in the clear supernatant was quantitated by
15 spectroscopic evaluation at a wavelength of 280 nm, using an
extinction coefficient of 2.015 for a 1 mg/ml solution of
V8-GLP-1 in a 1 cm cell. The crystallization yield was
quantitated by subtracting the V8-GLP-1 level in the
supernatant from the V8-GLP-1 level in the starting
20 solution. This sample showed a crystallization yield of
92.4%.

Example 16**pH 7.4 with 8.6% Ethanol and Zinc**

25 V8-GLP-1 was prepared at 4 mg/ml in 50 mM glycine
pH 9.0 buffer, followed by passage through a 0.2 μ m filter
(Gelman Sciences, Ann Arbor, Michigan). To 5 ml of this
solution was added 500 μ l of absolute ethanol followed by
300 μ l of a 2.08 mg/ml zinc chloride solution in water.
30 Small additions of 1N HCl were used to adjust the solution

-27-

to pH 7.40. After gentle swirling the final solution was enclosed in a 10-ml glass vial and stored at ambient temperature for two days. Microscopic examination revealed modest-sized, individual tetragonal crystals. The V8-GLP-1 content remaining in the clear supernatant was quantitated by spectroscopic evaluation at a wavelength of 280 nm, using an extinction coefficient of 2.015 for a 1 mg/ml solution of V8-GLP-1 in a 1 cm cell. The crystallization yield was quantitated by subtracting the V8-GLP-1 level in the supernatant from the V8-GLP-1 level in the starting solution. This sample showed a crystallization yield of 85.0%.

Example 17

pH 6.4 with 5% Ethanol and 1.0% Ammonium Sulfate

V8-GLP-1 (12.5 mg) was weighed into a 20 ml glass vial. 2.0 ml of a 10 mM ammonium acetate buffer containing 150 mM NaCl at pH 6.4 was added. The pH of the turbid solution was clarified by adjustment to pH 9.5 with 5N NaOH, then lowered to pH 6.4 with 2N HCl. The clear solution was filtered through a 0.22 μ m Millex GV 13 syringe filter (Millipore, Bedford, MA) into a new 20 ml glass vial. The concentration of the V8-GLP-1 stock solution was determined from the absorbance at 280 nm using an extinction coefficient of 2.015 for a 1.0 mg/ml solution of V8-GLP-1 in a 1 cm cell. The protein concentration was adjusted to 5.0 mg/ml. A pH 6.4 precipitant solution containing 10 mM ammonium acetate, 150 mM NaCl, 2% ammonium sulfate and 10% ethanol was prepared and filtered through a 0.22 μ m Millex GV13 syringe filter. 2 ml of the precipitant solution was

-28-

slowly added to 2 ml of the V8-GLP-1 stock solution in a glass vial. The vial was gently swirled and incubated at room temperature for 2 days. Tetragonal plate-shaped crystals were observed with a yield of 92%.

5 The pH of the crystal suspension was adjusted to pH 5.5 with 1N HCl and zinc chloride was added to a final concentration of 0.15 mg/ml. After zinc soaking overnight at room temperature, the pH of the suspension was adjusted to pH 7.5 with 1N NaOH and the preservative meta-cresol was
10 added to a concentration of 3.16 mg/ml. This example shows that, if desired, preserved formulations of GLP-1 crystals can be prepared directly for pharmaceutical use without isolation of the crystals by centrifugation or filtration in an intermediate step.

15

Example 18

pH 7.6 with 8.5% Ethanol and Zinc

V8-GLP-1 was prepared at 4 mg/ml in 50 mM glycine pH 9.0 buffer, followed by passage through a 0.2 μ m filter
20 (Acrodisc from Gelman Sciences, Ann Arbor, Michigan). To 10 ml of this solution was added 1 ml of absolute ethanol followed by 600 μ l of a 6.7 mg/ml zinc acetate (2-hydrate) solution in water. 100 μ l of 2% acetic acid was added, resulting in a pH of about 7.6. After gentle swirling the
25 final solution was enclosed in a 20-ml glass vial and stored at ambient temperature for 24 hours. Microscopic examination revealed modest-sized, individual tetragonal crystals. To the entire solution was then added 3.555 ml of a solution containing 3.5 ml of a 14 mg/ml solution of m-
30 cresol in water and 55 μ l of 2% acetic acid, resulting in a

-29-

suspension with a final pH of about 7.2. After gentle swirling the suspension was enclosed in a 20-ml glass vial and stored at ambient temperature for 24 hours. Microscopic examination again revealed modest-sized, individual
5 tetragonal crystals.

After centrifugation of an aliquot for 5 minutes at ambient temperature, the V8-GLP-1 content remaining in the clear supernatant was determined by HPLC analysis of a diluted sample compared to HPLC analysis of V8-GLP-1
10 standard solutions. The crystallization yield was quantitated by subtracting the V8-GLP-1 level in the supernatant from the V8-GLP-1 level in the starting solution. This preserved V8-GLP-1 formulation showed a crystallization yield of 97.7 %.

15

Example 19

Crystal Stability

Single tetragonal crystals of V8-GLP-1 were prepared in 10 mM NH_4OAc , 1% $(\text{NH}_4)_2\text{SO}_4$, 10% ethanol buffer at
20 pH 6.4 at 18 °C as described in Example 8. The crystals were removed from the mother liquor by low speed centrifugation and resuspended in a 10 mM NH_4OAc , 16 mg/ml glycerin, pH 5.5, buffer to a concentration of about 4.9 mg/ml of V8-GLP-1.

2 ml of this suspension was low speed centrifuged
25 and the supernatant was removed by pipette. The pellet was resuspended in 4 ml of a 10 mM ammonium acetate, 16 mg/ml glycerin pH 5.5 buffer containing 0.1 mg/ml zinc. This crystal suspension was allowed to soak in the zinc solution overnight at 4 °C

-30-

The zinc-soaked crystal suspension was divided into four 1-ml aliquots. These suspensions were low speed centrifuged and their supernatants were removed by pipette. Four crystal suspensions were prepared in 10 mM ammonium acetate, 16 mg/ml glycerin, 0.1 mg/ml zinc buffer at pH 6.0. Further pH adjustments to pH 7.4 with 0.1N NaOH and/or additions of meta-cresol to a final concentration of 3.16 mg/ml were made to selected samples as illustrated in Table 2. Each suspension was further divided in half for storage at both room temperature (about 22 °C) and at 4 °C, providing a total of 8 test samples as shown in Table 2.

After 10 days, the crystal suspensions were examined under the microscope. The suspension filtrates were then evaluated by HPLC to quantitate the soluble V8-GLP-1 in the crystalline suspensions. The HPLC results are reported in Table 2.

-31-

Table 2:
Soluble V8-GLP-1 in Crystal Suspensions
after Storage for 10 days.

Sample	pH	Storage Temperature	mg/ml meta-cresol	Soluble V8-GLP-1 by HPLC
A	6.0	4 °C	0	0.19 %
B	7.4	4 °C	0	0.10 %
C	6.0	4 °C	3.16	0.05 %
D	7.4	4 °C	3.16	0.06 %
E	6.0	22 °C	0	0.16 %
F	7.4	22 °C	0	0.08 %
G	6.0	22 °C	3.16	0.03 %
H	7.4	22 °C	3.16	0.04 %

5

This experiment showed that less than 0.2 % of the V8-GLP-1 peptide became solublized in either the preserved or non-preserved crystal formulations over a 10-day period.

Microscopically, the crystal suspensions showed less agglomeration or clumping of the single, tetragonal crystals at pH 7.4 than at pH 6.0, and less at 4 °C than at 22 °C. The meta-cresol did not seem to have a significant effect on crystal agglomeration. Additional testing showed the presence of more than 3% ethanol in the crystal suspension, either from the original crystallization mother liquor or from subsequent additions, greatly reduced the clumping tendency of the crystals in both preserved and non-preserved formulations. Further tests revealed that, although the crystals are relatively stable in the presence of meta-cresol, they are less stable in the presence of 5 %

20

-32-

phenol, which slowly leads to the formation of amorphous material.

Additional crystal stability tests showed that the V8-GLP-1 crystals prepared at pH 6.4 are very stable chemically, with no degradation peaks observed by HPLC analysis after storage at 5 °C or room temperature for up to two months.

Stability tests of crystals prepared in glycine buffer as described in Example 16 showed the V8-GLP-1 crystals stored in the original mother liquor were not stable when meta-cresol was added to the level of 3.16 mg/ml. This test resulted in dissolution of the crystals after only 1 day. The crystal instability in this composition could be effectively blocked by addition of zinc (via a zinc chloride solution) prior to addition of the preservative.

WE CLAIM:

1. A process for preparing single rod-shaped or plate-like crystals of a glucagon-like peptide-1 related molecule (GLP) which comprises preparing a crystallization solution comprising a GLP, a buffering agent, an alcohol or a mono or disaccharide, and optionally, ammonium sulfate or zinc.
2. The process of Claim 1 wherein the GLP is at a final concentration of between about 1-10 mg/ml and is selected from the group consisting of a GLP-1 analog, a GLP-1 derivative, a dipeptidyl-peptidase-IV (DPP-IV) protected GLP, a GLP-1 peptide analog, or a biosynthetic GLP-1 analog, and wherein the buffering agent is about 10 to 50 mM, and about pH 6-7 and is selected from the group consisting of Tris, ammonium acetate, sodium acetate, or Bis-Tris, and wherein the alcohol or mono or disaccharide is selected from the group consisting of methanol, ethanol, propanol, glycerol, trehalose, mannitol, glucose, erythrose, ribose, galactose, fructose, maltose, sucrose, and lactose, and, optionally, wherein about 1% ammonium sulfate is present.
3. The process of Claim 1 wherein total zinc is in a 0.5-1.7 molar ratio to the GLP which is at a final concentration of between about 1-20 mg/ml and is selected from the group consisting of a GLP-1 analog, a GLP-1 derivative, a DPP-IV protected GLP, a GLP-1 peptide analog, or a biosynthetic GLP-1 analog, and wherein the buffering agent is about 10 to 100 mM, and about pH 7-10 and is selected from the group consisting of glycine, aspartic acid

-34-

or Tris, and wherein the alcohol or mono or disaccharide is selected from the group consisting of methanol, ethanol, propanol, glycerol, trehalose, mannitol, glucose, erythrose, ribose, galactose, fructose, maltose, sucrose, and lactose.

5

4. The process of any one of Claims 1 to 3 wherein the GLP is selected from the group consisting of a DPP-IV protected GLP, or a biosynthetic GLP.

10

5. The process of any one of Claims 1 to 4 wherein the GLP is a DPP-IV protected GLP selected from the group consisting of Val-8-GLP-1(7-37)OH, Thr-8-GLP-1(7-37)OH, Gly-8-GLP-1(7-37)OH, or Met-8-GLP-1(7-37)OH.

15

6. The process of any one of Claims 1 to 5 having the additional step of soaking the GLP crystals in a zinc containing solution.

20

7. GLP crystals having tetragonal flat rod shaped or plate-like morphology selected from the group consisting of a GLP-1 analog, a GLP-1 derivative, a DPP-IV protected GLP, a GLP-1 peptide analog, or a biosynthetic GLP-1 analog.

25

8. The crystals of Claim 7 wherein the GLP is selected from the group consisting of DPP-IV protected GLP, or a biosynthetic GLP.

30

9. The crystals of Claim 7 wherein the GLP is selected from the group consisting of Val-8-GLP-1(7-37)OH, Thr-8-GLP-1(7-37)OH, Gly-8-GLP-1(7-37)OH, or Met-8-GLP-1(7-37)OH.

-35-

10. GLP crystals whenever prepared by the process of any one of Claims 1 to 6.

5 11. A substantially homogenous composition of GLP crystals.

12 The composition of Claim 11 wherein the GLP crystals are selected from the group consisting of a GLP-1 analog, a GLP-1 derivative, a DPP-IV protected GLP, a GLP-1 peptide analog, or a biosynthetic GLP-1 analog.

13. The composition of any one of Claims 11 to 12 wherein the GLP is selected from the group consisting of DPP-IV protected GLP, or a biosynthetic GLP.

14. The composition of any one of Claims 11 to 13 wherein the GLP is selected from the group consisting of Val-8-GLP-1(7-37)OH, Thr-8-GLP-1(7-37)OH, Gly-8-GLP-1(7-37)OH, or Met-8-GLP-1(7-37)OH.

15. A pharmaceutical formulation comprising a GLP crystal as claimed in any one of Claims 7 to 11 together with one or more pharmaceutically acceptable diluents, carriers or excipients therefor.

16. The pharmaceutical formulation of Claim 15 wherein the formulation is prepared by additions to and/or modifications of the post-crystallization mother liquor without separating the GLP crystals from the mother liquor.

-36-

17. A method of treating diabetes, obesity or related conditions in a mammal in need thereof, which comprises administering to said mammal a GLP crystal of any one of Claims 7 to 10.

5

18. A method of treating diabetes, obesity or related conditions in a mammal in need thereof, which comprises administering to said mammal a composition of any one of Claims 11 to 14.

10

19. A method of treating diabetes, obesity or related conditions in a mammal in need thereof, which comprises administering to said mammal a pharmaceutical formulation of any one of Claims 15 to 16.

15

20. The GLP crystals as claimed in any one of Claims 7 to 10 for use in treating diabetes, obesity or related conditions.

20

21. A substantially homogenous composition of GLP crystals as claimed in any one of Claims 11 to 14 for use in treating diabetes, obesity or related conditions.

25

22. A GLP crystal substantially as hereinbefore described with reference to any one of the Examples.

23. A substantially homogenous composition of GLP crystals substantially as hereinbefore described with reference to any one of the Examples.

30

SEQUENCE LISTING

<110> Eli Lilly and Company

<120> GLUCAGON-LIKE PEPTIDE-1 CRYSTALS

<130> X-10242 PCT

<140> PCT/US98/26480

<141> 1998-12-14

<150> US 60/069728

<151> 1997-12-16

<160> 4

<170> PatentIn Ver. 2.0

<210> 1

<211> 31

<212> PRT

<213> Homo sapiens

<400> 1

His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly
 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly
 20 25 30

<210> 2

<211> 29

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetically
 mutated human sequence

<220>

<223> Xaa at position 13 is Glu, Gln, Ala, Thr, Ser or
 Gly; Xaa at position 19 is Glu, Gln, Ala, Thr,
 Ser or Gly; and Xaa at position 29 is Gly or
 absent.

<400> 2

Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Xaa Gly Gln Ala

1 5 10

Ala Lys Xaa Phe Ile Ala Trp Leu Val Lys Gly Arg Xaa

 20 25

<210> 3
 <211> 29
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetically mutated human sequence.

<220>
 <223> Xaa at position 28 is Lys or absent; Xaa at position 29 is Gly or absent; and, if Xaa at position 28 is absent, Xaa at position 29 must be absent.

<400> 3
 His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly

 1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Xaa Xaa

 20 25

<210> 4
 <211> 30
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetically mutated human sequence

<220>
 <223> Xaa at position 19 is Lys or Arg; Xaa at position 30 is Gly or is absent; and Lys at position 27 may be acylated.

<400> 4
 Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln

 1 5 10 15

Ala Ala Xaa Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Xaa

20

25

30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/26480

A. CLASSIFICATION OF SUBJECT MATTER														
IPC(6) :A61K 38/26; C07K 14/605 US CL :514/12, 53, 866,909; 530/308, 324 According to International Patent Classification (IPC) or to both national classification and IPC														
B. FIELDS SEARCHED														
Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/12, 53, 866,909; 530/308, 324														
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched														
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Extra Sheet.														
C. DOCUMENTS CONSIDERED TO BE RELEVANT														
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
Y	EP 0 658 568 A1 (ELI LILLY AND COMPANY) 21 June 1995, page 4, lines 4 to page 7, lines 58.	1-4, 7-9, 11-13 and 22-23												
Y	EP 0 619 322 A2 (PFIZER INC.) 12 October 1994, entire document, especially page 18, lines 28 to page 19, lines 10.	1-4, 7-9, 11-13 and 22-23												
A,P	EP 0 869 135 A1 (ELI LILLY AND COMPANY) 07 October 1998, page 6, lines 36 to page 7, lines 26.	1-4, 7-9, 11-13 and 22-23												
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"> <tr> <td>* Special categories of cited documents:</td> <td>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>*A* document defining the general state of the art which is not considered to be of particular relevance</td> <td>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>*E* earlier document published on or after the international filing date</td> <td>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>*A* document member of the same patent family</td> </tr> <tr> <td>*O* document referring to an oral disclosure, use, exhibition or other means</td> <td></td> </tr> <tr> <td>*P* document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </table>			* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family	*O* document referring to an oral disclosure, use, exhibition or other means		*P* document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention													
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art													
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family													
O document referring to an oral disclosure, use, exhibition or other means														
P document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search 30 MARCH 1999		Date of mailing of the international search report 22 APR 1999												
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer ABDEL A. MOHAMED <i>Jas</i> Telephone No. (703) 308-0196												

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/26480

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PRIDAL et al. Absorption of Glucagon-like Peptide-1 can be Protected by Zinc or Protamine. 1996, Vol. 136, No. 1-2, pages 53-59, especially page 53.	1-4, 7-9, 11-13 and 22-23

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/26480

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 5, 6, 10 and 14-21
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

- Remark on Protest The additional search fees were accompanied by the applicant's protest.
 No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/26480

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, CAS ONLINE, MEDLINE, BIOSIS, WPIDS, EMBASE

search terms: GLP# or GLUCAGON(1W)PEPTIDE OR (DPP(W) (TV OR 4) OR DPP4) (S) PEPTIDASE OR
(DIPEPTIDYL OR DI PEPTIDYL) (W) PEPTIDAS) (S) (MANUF? OR PROD? OR PREP?); BUFFER? OR TRIS OR
AMMON? OR NA OR SODIUM (W) ACETATE; MONOSACCHARID? OR DISACCHARIDE? OR METANOL OR
ETH? OR ALCOHOL; TREAT? OR THERAPEUT? DIABETES OR OBESITY; CRYSTAL? OR
PHARMACEUT?(W)COMPOSIT?